

Possible scenarios that the "New Horizons" spacecraft may find in its close encounter with Pluto

Héctor Javier Durand-Manterola and Héctor Pérez-de-Tejada
Instituto de Geofísica, Universidad Nacional Autónoma de México (UNAM)
hdurand_manterola@yahoo.com

Abstract

Next year (2015) the "New Horizons" spacecraft will have a close encounter with Pluto. In the present study we discuss some possibilities regarding what the spacecraft may encounter during its approach to Pluto. Among them we should include: the presence of geological activity due to heat generated by tides; the unlikely presence of an intrinsic magnetic field; the possibility of a plasmasphere and a plasmopause; the position of an ionopause; the existence of an ionospheric terminator flow similar to that at Venus and Mars; and the presence of a Magnus force that produces a deflection of Pluto's plasma wake. This deflection oscillates up and down in its orbit around the sun.

Key words: Pluto, Solar Wind Interaction, Magnus effect

1 Introduction

In 1976 frozen CH₄ was identified in the surface of Pluto (Cruikshank et al., 1976; Hansen y Paige, 1996). It was known that Pluto has a bright surface, and since methane ice tends to darken when subjected to radiation, it was proposed that sublimation and transport of methane keep fresh the surface (Stern et al., 1988), thus implying the presence of an atmosphere. Through stellar occultation a tenuous atmosphere was detected in 1988, and assuming that nitrogen was its main constituent, a pressure between 0.2 and 0.5 Pa was calculated (Elliot et al., 1989; Hubbard et al., 1990; Elliot y Young, 1991; Hansen y Paige, 1996). Finally, in 1992 the spectral signatures of nitrogen and carbon monoxide were identified, the two species occurring in solid state (Owen et al., 1992; Hansen y Paige, 1996).

Comparisons have been made between Triton and Pluto since both bodies have similar density, radius, rotation speed, surface gravity, atmospheric pressure and albedo. The volatile inventory of both, include N₂, CH₄ y CO. Both bodies have a thin atmosphere whose main component is nitrogen. Both have bright poles (Hansen y Paige, 1996).

The nature of the interaction of the solar wind with the ionosphere of Pluto depends on the amount of material the atmosphere lost. If Pluto has a large atmospheric escape

flux interaction will be comet type. On the other hand if it has a weak atmospheric escape flux interaction will be Venus type (Bagenal and McNutt, 1989)

The New Horizons spacecraft will reach Pluto in July 2015 and will be the first approach to a dwarf planet in the Kuiper belt and the second visit to a dwarf planet after the visit of the "Dawn spacecraft" to Ceres in February of that year. In this study, we explore some possibilities about what the "New Horizons may find in its approach to Pluto.

2 Scenarios

2.1 Intrinsic Magnetic field in Pluto

If Pluto has a fluid inner layer and this layer is conductive, then Pluto may have a dynamo that generates a magnetic field. Durand-Manterola (2009) obtained the following empirical equation, which relates the mass of the planet (M_p), the conductivity of the conductive layer (σ) and the period of rotation of the body (P_e), with the planetary magnetic moment (M):

$$M = 1 \times 10^{-5} \left[\frac{M_p \sigma}{P_e} \right]^{1.106} \quad (2.1.1)$$

Using this equation we can calculate the magnetic field present at Pluto. If we assume that Pluto has a (very unlikely) molten iron core, then with the appropriate values (Table 1), the equation (2.1.1) gives the following value of the magnetic moment: 5.34×10^{18} Amp-m², which is one order of magnitude smaller than the magnetic moment of Mercury.

Pluto is like the icy moons of the giant planets. Of these, Ganymede is the only with an intrinsic magnetic field. It has been suggested that Ganymede's magnetic field is generated in an underground saltwater ocean (Kivelson et al., 2002). If this is also the case for Pluto its conductivity will be less than that assuming iron. With the maximum value of seawater conductivity (Table 1) equation (2.1.1) gives a magnetic moment of 1.03×10^{14} Amp-m².

If Pluto has an intrinsic magnetic field, then it has a magnetopause, which in the subsolar point, will be at the distance (Eq. (12) in Durand-Manterola, 2009, corrected):

$$R_{ss} = \left[\frac{\mu_0 M^2}{32\pi^2 \rho V^2} \right]^{1/6} \quad (2.1.2)$$

where M is the dipolar magnetic moment, μ_0 is the permeability of free space and ρ and V are the density and velocity of the solar wind.

Applying the appropriate values (Table 1) in this equation and taking into account an iron core (which could be very unlikely), the subsolar point of the magnetopause will be 5768 km from the centre of Pluto, i.e., 4.95 Plutonian radii.

If the magnetic field is generated by an ocean of saltwater, equation (2.1.2) leads to $R_{ss} \sim 161.7$ km which is an impossible value since it is smaller than Pluto's radius. In this case the magnetopause would be by the surface of Pluto or at the ionopause. Under such circumstances the presence of a magnetic field could only be verified in the wake where the magnetospheric plasma is not compressed.

About 50 minutes after its closest approach to Pluto the "New Horizons" will move through its wake. However, the spacecraft does not have a magnetometer and thus a magnetopause only will be inferred from observations of a discontinuity in the plasma properties such as a jump in density. On Earth the density and the temperature of the magnetospheric plasma gradually decrease within the magnetotail, being the highest at the magnetopause (Rosenbauer et al., 1975). If Pluto's magnetosphere is similar to that at Earth, we should expect such variations in density and temperature.

Before reaching the magnetopause, the solar wind will be influenced by a shock wave, and thus equation (2) will not be applicable since it involves a balance between the internal magnetic force and the kinetic energy of the free solar wind. However, when the shock wave slows down the solar wind it is compressed together with the convected magnetic field thus increasing the magnetic pressure (the kinetic pressure is converted into magnetic pressure). Therefore, a pressure balance is set between the magnetic pressure of the solar wind and the magnetic pressure of the planet. As long as the magnetic pressure of the solar wind is equal to the kinetic pressure, then (2.1.2) is applicable.

2.1.1 Plasmasphere and plasmopause

Inside the magnetospheres of planets there is a plasma zone corotating with the planet, the plasmasphere, in which the plasma provided by the ionosphere is driven by the corotation electric field. The distance to the boundary of this region, or plasmopause can be calculated by equating the centripetal force to the sum of the gravitational and the convective forces. From this we obtain:

$$r = \left[\frac{q\mu_0 M}{4\pi\omega m} + \frac{GM_p}{\omega^2} \right]^{1/3} \quad (2.1.3)$$

Assuming that there are only protons in the Plutonian plasmasphere and using the values of Table 1 we find that the corotation zone has a radius of 19,100 km, for the lower magnetic moment; and 238,000 km for the larger magnetic moment. In either case the radius of the corotation zone is greater than the distance to the subsolar magnetopause. Therefore, the whole frontal magnetosphere is corotating with Pluto.

For Pluto has a magnetic field requires that inside there is a conductive liquid layer. For this to happen requires an internal heat source. One possible source would tidal heating.

2.2 Tidal heating of the planetary body

Pluto and its moon Charon have trapped rotation, i.e., both bodies show each other the same hemisphere. On the other hand the orbit of Charon is circular (and therefore so is Pluto's orbit around their common centre of mass) (Buie et al., 2012). For these two conditions it is not expected that there is an exchange of energy, via tides, between the two bodies.

However, solar gravity force modifies this view since Charon may be in conjunction, in opposition, square and all intermediate positions with respect the Sun, thus both bodies will exert on Pluto a continuous changing tidal force that implies a continuous, and periodic, deformation of Pluto's body (and of course Charon's body). This continuous exchange of energy in tidal forces produces heat due to dissipation of tidal energy. The heat flow may manifest itself in cryo-volcanic activity on the surfaces of both bodies, or in the presence of a molten layer in their interior, or both. This happens only in certain positions in the orbit of Pluto around the Sun (see Figure 1). As the axis of rotation of the planet is lying almost in the plane of its orbit then the plane of the orbit of Charon is almost perpendicular to the orbital plane. As this plane is fixed respect to the background stars this means that there will be times when the plane of the orbit of Charon will be perpendicular to the direction Sun-Pluto and at other times will be parallel. In the latter situation would be where would have tidal effects.

If F_C is the gravitational force exerted on Pluto by Charon, and F_S is the gravitational force exerted by the Sun on Pluto, then the ratio of F_C to F_S is:

$$\frac{F_C}{F_S} = \frac{M_C}{M_S} \frac{R_S^2}{R_C^2} \quad (2.2.1)$$

where M are the masses and R the distances of the Sun and Charon to Pluto and the subscripts S and C correspond, respectively, to both bodies. Putting the appropriate values (Table 1) in equation (2.2.1) we obtain that the force exerted by Charon on Pluto is 39.3 times the force exerted by the Sun in the perihelion and 108.4 times in the aphelion.

At perihelion, when the Sun and Charon are in conjunction as seen from Pluto the force exerted will be the sum of the two forces, i.e. 40.3 times the force exerted by the Sun. Three days later when Charon is in opposition to the Sun, the force exerted on Pluto is the difference between both forces, i.e., 38.3 times the force applied by the Sun. The difference between both positions is $\sim 5\%$. At aphelion, during the conjunction, the sum of the Sun and Charon forces is 109.4 times the force of the Sun, and in opposition 107.4 times (This difference is 1.8%). The heat released by the deformation produced by these differential forces may aid in the sublimation of N_2

and CH_4 , which are solid at the surface, and thus can help the formation of the atmosphere. Moreover, since Charon is less massive than Pluto, such dissipation may explain the fact that the surface of Charon is warmer than Pluto (Marcialis et al., 1987).

2.3 Ionopause on Pluto

The temperature on Pluto's surface is 40 ± 2 °K (Strobel et al., 1996). Given that the atmosphere is formed mainly of N_2 , then the thermal velocity is 150 m/s. According to the model of Strobel et al., (1996) the atmospheric temperature at 80 km altitude reaches between 100 y 130 °K. Taking these values, the thermal velocities are 240 m/s and 280 m/s, respectively. On the other hand, since the escape velocity on Pluto is 1220 m/s (Tholen et al., 2000) the thermal velocities have smaller values and thus Pluto's atmosphere remains trapped.

Ions were detected in the Plutonian atmosphere (Young et al., 2008), that is, Pluto has an ionosphere. For all the above, and in the case that Pluto does not have a magnetic field or that it has a weak magnetic field generated by an ocean of salt water, then there will be a ionopause. We can calculate the position of the ionopause with a pressure balance between the thermodynamic pressure of the ionospheric plasma and the ram pressure of the solar wind.

$$\rho V^2 \cos^2 \xi = n_i k T \quad (2.3.1)$$

Where ρ is the density of the solar wind, V is the solar wind speed in free space at the distance of Pluto, ξ is the subsolar angle, n_i and T are the particle density and the ionospheric plasma temperature, and k is the Boltzmann constant.

If the solar wind arrives perpendicular to the ionosphere, as in the subsolar point ($\xi = 0$), then all its momentum is used to compress the ionosphere and ionopause be formed where the pressure and the thermal pressure of the ionosphere are balanced. But if the wind gets tilted to the vertical at some point in the ionosphere ($\xi \neq 0$) then its momentum will have two components: a vertical which will compress the ionosphere, and an horizontal which will have no effect on it. In this case the wind ram pressure will be lower, then the pressure of the ionosphere is balanced at higher altitudes where the density is less.

Before reaching the ionosphere, the solar wind decelerates as it moves across the shock wave; then strictly the pressure balance will occur between the magnetic field pressure plus the ram pressure of the shocked solar wind and the thermal pressure of the ionosphere. Since the magnetic pressure and the shocked ram pressure are equivalent to the free stream solar wind kinetic pressure then equation (2.3.1) is applicable.

From equation (2.3.1) and solving for n_i , we have:

$$n_i = \frac{n_{VS} m_P V^2}{kT} \cos^2 \xi \quad (2.3.2)$$

Thus leading to the particle density of the ionosphere at the ionopause.

Moreover, from the model of Ip et al. (2000) we can obtain the altitude corresponding to the particle density value; namely:

$$h = 2.2611 \times 10^3 \exp(-3 \times 10^{-4} n_i) \quad (2.3.3)$$

Combining equations (2.3.2) and (2.3.3) lead to the ionopause altitude. Figure (2.3.1) shows values of this altitude as a function of solar zenith angle. The altitude at the subsolar point is 2147 km and 2261 km by the terminator, or measured from the planet centre, 3312 km at the subsolar point, and 3426 km by the terminator.

When the Sun-Pluto line is in the plane of Charon's orbit the solar wind may not reach the ionopause by the subsolar point thus allowing the Plutonian ionosphere to expand.

2.4 Trans-terminator flow

When there is a magneto-hydrodynamic interaction between the solar wind flow and Pluto's ionosphere (basically wave-particle interactions), there should be momentum transfer from the solar wind to the upper ionosphere, which generates a flow of ions from dayside to night side (Perez-de-Tejada, 1986). This type of flow was first discovered at Venus and is called the ionospheric trans-terminator flow (Knudsen et al., 1980). A similar trans-terminator flow is expected to occur at Pluto. We can calculate its speed from an equation that relates the conservation of momentum flux between the solar wind and the ionosphere, i.e.:

$$\pi r_i^2 \rho_{SW} V_{SW}^2 = \pi (r_i^2 - r_0^2) \rho_i V_i^2 \quad (2.4.1)$$

Here $\rho_{SW} V_{SW}^2$ is the momentum flux density of the solar wind and $\rho_i V_i^2$ is the momentum flux density in the ionosphere, r_i is the radius of the ionopause at the terminator and r_0 is the radius of Pluto. Solving for V_i we have:

$$V_i = V_{SW} \left[\frac{r_i^2}{r_i^2 - r_0^2} \frac{\rho_{SW}}{\rho_i} \right]^{1/2} \quad (2.4.2)$$

Using the values of Table 1 we obtain 2.8 km/s, which is larger than the escape velocity of 1.22 km/s and thus the momentum transfer from the solar wind to the Pluto ionosphere should produce a significant loss of ionospheric material.

2.5 Magnus force at Pluto

When a sphere moves through a fluid and at the same time rotates, a dynamic force, is produced and is oriented away from the directional flow (Magnus, 1853). Since Pluto is also immersed in the solar wind and is a rotating sphere it could also produce a Magnus force. However, since the solar wind is non-collisional plasma, it could interact directly with Pluto's solid body and thus the Magnus force, which is a hydrodynamic effect, would not occur. However, since Pluto has an ionosphere or a plasmasphere, its interaction with the solar wind could produce a magneto-hydrodynamic interaction resulting in a Magnus force. If Pluto has an intrinsic magnetic field the plasma of all the magnetosphere co-rotates with the planet, as we see in section 2.1.1. If it does not have a magnetic field then the lower ionosphere co-rotates with the solid body and the high region flows from dayside to night side (trans-terminator flow). In this case the Magnus effect occurs between the trans-terminator flow and the internal rotating ionosphere, but the effect of deviation of the wake, is manifests even in the solar wind.

Previous work has suggested that the Magnus force has an effect on the plasma wake of Venus and Mars (Perez-de-Tejada, 2006; Perez-de-Tejada et al., 2009). Such force was used to account for the deviated direction of the Venus plasma wake as a result of the solar wind interaction with the Venus ionosphere (Pérez-de-Tejada, 2006, 2008). It is possible that similar conditions are also applicable to Pluto's wake.

Since the plane of the equator of Pluto has 119.61° inclined to the plane of the orbit (Tholen et al., 2000) then as it moves in its orbit around the sun there will be times when the rotation axis is perpendicular to the solar wind flow. At this time is when will be Magnus effect. The Magnus effect will produce a deviation of the wake of the planet relative to the direction Sun-Pluto. In other parts of the orbit, when the rotation axis is parallel to the solar wind, there is no Magnus effect and the wake will be aligned with the Sun-planet line. In some parts of the Pluto's orbit the Magnus force points to the solar north and in others it points to the solar south (Pérez-de-Tejada et al., 2014). Such change will produce a cyclic variation of the wake deviation along Pluto's orbit. At times it will be diverted to the north of the orbital plane and in others to the south.

The Magnus force applied to a rotating planet is:

$$f_M = 4\pi\rho V\omega R^3 \quad (2.5.1)$$

Where ρ and V are the density and speed of the solar wind, and ω is the angular velocity of rotation of the planet. R is the radius of the corotating plasma (ionopause or magnetopause) but not the radius of the planet since the Magnus force arises from the magneto-hydrodynamic interaction between the solar wind and a planetary plasma (ionosphere or plasmasphere).

2.6 Angle of the wake

The geometry of the flow generated by the Magnus force causes a deflection angle of the plasma wake. If the rotation speed of Pluto's ionosphere (or plasmasphere) is V_P

and the velocity of the flow is V_f then in the upper ionosphere (or in the magnetopause) where both velocities are in the same direction the total speed is:

$$V_1 = V_f + V_p \quad (2.6.1)$$

And in the side where the two velocities are anti-parallel we have:

$$V_2 = V_f - V_p \quad (2.6.2)$$

Consider two parcels of flow together in the sub-solar point. One of the parcels moves to the side where the rotation motion and the flow are parallel and the other moves where the flow and the rotation motion are anti-parallel. The positions with time of the two parcels around Pluto are:

$$x_1 = V_1 t_1 - \pi R = (V_f + V_p)t_1 - \pi R \quad (2.6.3)$$

$$x_2 = V_2 t_2 + \pi R = -(V_f - V_p)t_2 + \pi R \quad (2.6.4)$$

Where R is the radius of the corotating sphere (ionosphere or plasmasphere), and t_1 and t_2 are the travel times of parcels 1 and 2 in their trajectories. We take the coordinate x measured on the surface of the co-rotating sphere, and zero in the anti-solar point. Therefore, parcel 1 begins at $x_1 = -\pi R$ and the x -coordinate increases. Meanwhile, parcel 2 starts at $x_2 = \pi R$ and the x -coordinate decreases.

When the parcels encounter again, after moving around Pluto, their travel time is: $t_1 = t_2 = t$ and their position is: $x_1 = x_2 = x$. Then we put these values into equation (2.6.4) and solving for t we obtain:

$$t = \frac{x - \pi R}{V_p - V_f} \quad (2.6.5)$$

Substituting this value in equation (2.6.3), and after some algebraic manipulation, we have:

$$x = \pi R \frac{V_p}{V_f} \quad (2.6.6)$$

This is the distance where the two parcels encounter, measured from the anti-solar point. At this point the parcels separate from Pluto and form the plasma wake. If we divide the equation (2.6.6) by R we have the angle of the wake (in radians) from the Sun-Pluto direction:

$$\theta = \pi \frac{V_p}{V_f} \quad (2.6.7)$$

Multiplying by $180/\pi$ we have this angle in degrees:

$$\theta = 180 \frac{V_p}{V_f} \quad (2.6.7)$$

Due to the velocity of Pluto in its orbit around the sun the solar wind does not reach exactly on the Sun-Planet direction, but at an angle to the right (as seen from the Sun):

$$\varphi = \arctg\left(\frac{V_o}{V}\right) \quad (2.6.8)$$

Where V_o is Pluto's orbital velocity and V is the free stream solar wind velocity. Due to the fact that the planes in which the angles θ and φ occur are almost perpendicular to each other they cannot be added together. While φ keeps nearly a steady value ($\varphi = 0.6^\circ$ for Pluto's mean orbital velocity) the angle θ varies depending on the position occupied by Pluto in its orbit around the sun. Taking values from Table 1 we obtain a maximum $\theta = 1^\circ$ value if Pluto has an ionopause, or $\theta = 0.3^\circ$ if it has a magnetosphere.

3 Discussion and conclusions

The main scenarios that the New Horizons spacecraft may find at Pluto are:

3.1 Tidal heating. During perihelion differential tides exerted by Sun and Charon on Pluto is 5.1 % the force exerted by Charon. At aphelion it is 1.8 %. Such difference subjects the planetary body to tension and release heat that can be manifested as some kind of geological activity.

3.2 Magnetic field. Even though the presence of an intrinsic magnetic field in Pluto is unlikely it is still possible. If we assume that Pluto has a molten iron core (quite unlikely) then we can suggest that the magnetic moment is 5.34×10^{18} Amp-m², an order of magnitude smaller than that of Mercury. In case that an underground saltwater ocean produces a magnetic field, as in the case of Ganymede, the magnetic moment would be 1.03×10^{14} Amp-m². With the largest magnetic moment, the subsolar point of the magnetopause will be 5768 km from the center of Pluto, i.e., 4.95 Plutonian radius. With the smaller magnetic moment it would be by the ionopause.

3.3 Plasmasphere and Plasmopause. Assuming there are only protons in the Plutonian plasmasphere corotation zone has a radius of 19,100 km, for the lower magnetic moment, and 238,000 km for the higher magnetic moment. In either case the radius of corotation zone is larger than the distance to the sub-solar magnetopause, and thus the entire front of the magnetosphere co-rotates with the planet.

3.4 Ionopause. The height of the ionopause at the subsolar point is 2147 km and grows to 2261 km by the terminator. Measured from the center of the planet it is 3312 km, at the subsolar point, and 3426 km by the terminator.

3.5 Trans-terminator flow. The trans-terminator flow speed obtained from the model is 2.8 km/s, which is larger than the 1.22 km/s escape velocity. Therefore, the momentum transfer from the solar wind to the Pluto's ionosphere should produce a significant loss of ionospheric material.

3.6 Angle of the wake. Pluto's plasma wake is subject to two different deviations: A) the aberration angle of arrival of the solar wind since Pluto's orbital motion is perpendicular to the radial direction of flow (at Pluto the mean angle is $\varphi = 0.6^\circ$). B) The angle due to the Magnus force. At Pluto this angle is $\theta = 11.5^\circ$ if Pluto has an ionopause, or $\theta = 0.3^\circ$ if it has a magnetosphere. These are maximum values. Unfortunately, when the New Horizons spacecraft approaches Pluto, the conditions will not be optimal for observing a deviation of the Magnus angle in the wake because the solar wind direction and the rotation axis will be parallel. At Venus the aberration angle and the Magnus angle are in the same plane and thus they add up so that it is difficult to quantify the contribution of each deviation on the wake. In Pluto both angles are in nearly perpendicular planes, and thus they can be distinguished from each other.

References

- Bagenal, Fran and R.L. McNutt Jr. (1989). Pluto's interaction with the solar wind. *Geophys. Res. Lett.* 16(11): 1229-1232.
- Buie, M.W., D. J. Tholen, and W. M. Grundy. (2012). The orbit of charon is circular. *The Astronomical Journal*, 144:15 (19pp), 2012 July doi:10.1088/0004-6256/144/1/15
- Cox, A.N. (2000). Chapter 2 General Constants and Units. In *Allen's Astrophysical Quantities*. 4th Edition Editor A.N. Cox. AIP Press and Springer.
- Cruikshank, D. P., Pilcher, C. B. And Morrison, D. (1976). Pluto - Evidence for methane frost. *Science*, vol. 194, Nov. 19, 1976, p. 835-837.
- Delamere, P.A. and F. Bagenal. (2004). Pluto's kinetic interaction with the solar wind. *Geophysical Research Letters*, 31: L04807, doi:10.1029/2003GL018122
- Durand-Manterola, H.J. (2009). Dipolar magnetic moment of the bodies of the solar system and the Hot Jupiters. *Planet. and Space Sci.* 57: 1405-1411. (Also in arxiv: 1007.4497).
- Elliot, J.L. E.W. Dunham, A.S. Bosh, S.M. Slivan, and L.A. Young (1989). Pluto's radius and atmosphere: Results from the entire 9 June 1988 occultation data set. *Icarus* 77: 148-170.
- Elliot, J.L., and L.A. Young. (1992). Analysis of stellar occultation data for planetary atmospheres. I Model fitting with application to Pluto. *Astrom J.* 103: 991- 1015.
- Hansen, C.J. and D.A. Paige. (1996). Seasonal Nitrogen cycles on Pluto. *Icarus* 120: 247-265.
- Hubbard, W.B., R.V. Yelle, and J.I. Lunine. (1990). Nonisothermar Pluto atmosphere models. *Icarus* 84, 1-11.

- Ip, W.H., A. Kopp, L.M. Lara, and R. Rodrigo. (2000). Pluto's Ionospheric Models and Solar Wind Interaction. *Adv. Space Res.* Vol. 26, No. 10, pp. 1559-1563.
- Kanehl, P. (2010). Particle model of the Magnus effect. Bachelor Thesis. Mathematisch-Naturwissenschaftliche Fakultät Ernst-Moritz-Arndt-Universität Greifswald
- Kennish, M.J., 2001. *Practical Handbook of Marine Science*. CRC Press, Boca Raton, FL.
- Kharlamov, A., Z. Chara, and P. Vlasak. (2007). Magnus And Drag Forces Acting On Golf Ball. *Colloquium FLUID DYNAMICS 2007 Institute of Thermomechanics AS CR, v. v. i., Prague, October 24 - 26, 2007*.
- Kittel, C. (2004). *Introduction to Solid State Physics*. John Wiley & Sons, Inc. New York. Versión en español (1981) Editorial Reverte, Barcelona.
- Kivelson, M.G., K.K. Khuruna, M. Volwerk. (2002). The permanent and inductive magnetic moments of Ganymede. *Icarus* 157: 507-522.
- Knudsen, W.C., K. Spenner, K. L. Miller and V. Novak. (1980). Transport of ionospheric O⁺ ions across the Venus terminator and implications. *Journal of Geophysical Research: Space Physics* 85 (A13): 7803-7810
- Magnus, G. (1853). Über die Abweichung der Geschisse, und: Über eine abfallende Erscheinung bei rotierenden Körperen (On the deviation of projectiles, and: On a sinking phenomenon among rotating bodies). *Annalen der Physik*, vol 164, no 1, pages 1-29.
- Marcialis, R.L., G.H. Rieke, L.A. Lebofsky. (1987). The Surface Composition of Charon: Tentative Identification of Water Ice. *Science, New Series* 237(4820): 1349-1351 (sep. 11, 1987).
- Owen, T.C., T. Geballe, C. De Bergh, L.Young, J. Elliot, D. Cruikshank, T. Roush, B. Schmitt, R.H. Brown, and J.Green. (1992). Detection of nitrogen and carbón monoxide on the surface of Pluto. *Bull. Am. Astron. Soc.* 24: 961.
- Pérez-de-Tejada. (1986). Fluid dynamic constraints of the Venus ionospheric flow. *Journal of Geophysical Research: Space Physics* 91(A6): 6765-6770.
- Pérez-de-Tejada, H., (2006) Magnus force in the Venus ionosphere. *Journal of Geophys. Res., Geophys. Res.*, 111, A111105.
- Pérez-de-Tejada, H., Viscous magnus force for the rotating Venus ionosphere. *Astrophys. J.*, 2008, 676, L65-L68.
- Pérez-de-Tejada, H. And R. Lundin. (2009). Magnus Force Effects in the Rotating Ionospheres of Venus and Mars. *American Geophysical Union, Fall Meeting 2009*, abstract #P11B-1208.
- Pérez-de-Tejada, H., H. Durand-Manterola, M. Reyes-Ruiz, and R. Lundin, Pluto's plasma wake driven away from the ecliptic plane, *ICARUS* (In PRESS, 2014).
- Resnick, D. And D. Halliday. (1970). *Física (Parte II)*. Compañía Editorial Continental S.A.
- Rosenbauer, H., H. Grünwalt, M.D. Montgomery, G. Paschman, and N. Scopke. (1975). Heos 2 Plasma observations in the distant Polar Magnetosphere. *The Plasma Mantle. J. Geophys. Res.* 80(19): 2723-2737
- Stern, S.A., L.M. Trafton, G.R. Gladstone. (1988). Why is Pluto bright? Implications of the albedo and lightcurve behavior of Pluto. *Icarus* 75: 485-498.
- Strobel, D.F., X. Zhu, M. E. Summers and M. H. Stevens. (1996). On the Vertical Thermal Structure of Pluto's Atmosphere. *Icarus* 120, 266-289.

Tholen, D.J., V.G. Teifel and A.N. Cox. 2000. Chapter 12 Planets and Satellites. In Allen's Astrophysical Quantities. 4th Edition Editor A.N. Cox. AIP Press and Springer.

Young, L.A., S. Alan Stern, Harold A. Weaver, Fran Bagenal, Richard P. Binzel, Bonnie Buratti, Andrew F. Cheng, Dale Cruikshank, G. Randall Gladstone, William M. Grundy, David P. Hinson, Mihaly Horanyi, Donald E. Jennings, Ivan R. Linscott, David J. McComas, William B. McKinnon, Ralph McNutt, Jeffery M. Moore, Scott Murchie, Carolyn C. Porco, Harold Reitsema, Dennis C. Reuter, John R. Spencer, David C. Slater, Darrell Strobel, Michael E. Summers, G. Leonard Tyler. (2008). New Horizons: Anticipated Scientific Investigations at the Pluto System. Submitted to Space Science Review. 140(1-4): 93-127.




Figure 1. At point A the Sun and Charon are not aligned with Pluto, then there are no tidal forces. But at point B are aligned with Pluto at the conjunction and opposition, at that time there are tidal force.

Table 1
Values of the variables used in calculations

Variable	Value	Reference
Proton charge (q)	4.803×10^{-19} C	Cox, 2000
Gravity constant (G)	6.672×10^{-11} m ³ kg ⁻¹ s ⁻²	Cox, 2000
Solar Wind density (ρ)	1.67×10^{-23} kg/m ³	Delamere et al., 2004 ¹
Pluto-Charon distance (Rc)	19571.4 ± 4.0 km	Young et al., 2007
Seawater conductivity (max.)	6.5683 S/m	Kennish, 2001
Conductivity of iron	1.2×10^5 S/m	Kittel, 2004
Pluto-Sun distance (Rs) (aphelion)	7.37×10^9 km	Tholen et al., 2000 ¹
Pluto-Sun distance (Rs) (perihelion)	4.44×10^9 km	Tholen et al., 2000 ¹
Charon Mass (Mc)	1.52×10^{21} kg	Young et al., 2007
Proton Mass (m)	1.672×10^{-27} kg	Cox, 2000
Pluto Mass (Mp)	$1.305 \pm 0.006 \times 10^{22}$ kg	Young et al., 2008
Sun Mass (Ms)	1.989×10^{30} kg	Cox, 2000
Pluto rotation period (Pe)	6.38 days = 5.512×10^5 s	Buie et al., 2012
Permeability of vacuum (μ_0)	1.26×10^{-6} H/m	Resnick, 1970
Pluto radius (r ₀)	1165 ± 25 km	Young et al., 2008
Ionosphere temperature (T)	1000 K	Ip et al., 2000
Angular velocity of Pluto (ω)	1.14×10^{-5} rad/s	Buie et al., 2012 ¹
Solar Wind velocity (V)	4.5×10^5 m/s	Delamere et al., 2004
Orbital velocity of Pluto (V _o)	4.749 km/s	Tholen et al., 2000

¹ Calculation based on data from the authors cited.

Appendix

Meaning of the symbols used

f_M \equiv Magnus force.

F_C \equiv Gravity force of Charon on Pluto.

F_S \equiv Gravity force of the Sun on Pluto.

G \equiv Gravity Constant

h \equiv Height above the surface of Pluto.

k \equiv Boltzmann Constant.

M \equiv Magnetic Moment

M_p \equiv Pluto Mass

M_C \equiv Charon Mass

M_S \equiv Sun Mass

m \equiv Plasmasphere Particles Mass

m_p \equiv Proton Mass

n_i \equiv Ionosphere Particle Density

n_{sw} \equiv Solar Wind Particle Density

P_e \equiv Planet Rotation Period

q \equiv Proton Charge

r \equiv Corotation region radius

r_0 \equiv Pluto Radius

r_i \equiv Ionopause radius at terminator

R \equiv Ionopause or Magnetopause Radius

R_C \equiv Distance of Charon to Pluto

R_S \equiv Distance of the Sun to Pluto

R_{SS} \equiv Distance from the centre of Pluto to the magnetopause in the sub-solar point

t \equiv time

T \equiv Ionosphere Temperature

v_p \equiv Rotational velocity of Pluto's equator

v_0 \equiv Orbital Velocity of Pluto

V \equiv Solar Wind Velocity before the shock wave

V_f \equiv Velocity of the flow in the Magnus force.

V_i \equiv Trans-terminator flow velocity

V_p \equiv Ionosphere or plasmasphere rotation velocity

V_{sw} \equiv Shocked Solar Wind speed in the terminator ($\sim 1/8$ to $1/10$ of V)

V_1 \equiv Sum of the solar wind and rotation velocities of the ionosphere or plasmasphere

V_2 \equiv Difference of the solar wind and rotation velocities of the ionosphere or plasmasphere

x_1 \equiv Distance travelled by the solar wind, near the ionopause, on the side parallel to rotation of the ionosphere

x_2 \equiv Distance travelled by the solar wind, near the ionopause, on the side anti-parallel to rotation of the ionosphere

φ = Aberration angle of the solar wind

μ_0 = Vacuum Permeability

θ = Deflection angle of the wake due to the Magnus force

ρ = Solar Wind Density

ρ_i = Ionosphere Density

ρ_{sw} = Shocked Solar Wind Density (~ 8 to 10 times ρ)

σ = Fluid layer Conductivity

ξ = Subsolar angle

ω = Pluto Angular velocity